

# R3 Report

## REQUIREMENT, RATIONALE, REFERENCE

A complimentary publication of Joint Commission

Issue 16, June 25, 2018

*Published for Joint Commission-accredited organizations and interested health care professionals, R3 Report provides the rationale and references that Joint Commission employs in the development of new requirements. While the standards manuals also may provide a rationale, R3 Report goes into more depth. The references provide the evidence that supports the requirement. R3 Report may be reproduced if credited to Joint Commission. Sign up for [email](#) delivery.*

### Pain assessment and management standards for office-based surgeries

Effective Jan. 1, 2019, new and revised pain assessment and management standards will be applicable to all Joint Commission-accredited ambulatory care, critical access hospitals, and office-based surgery organizations. Joint Commission published new and revised pain assessment and management requirements in January 2018 for the Hospital Accreditation program (see July 2017 issue of Perspectives). This project is a continuation of this initiative.

### Engagement with stakeholders, customers, and experts

In addition to an extensive literature review and public field review, Joint Commission sought expert guidance through the following:

- [Technical advisory panel \(TAP\)](#) included practicing clinicians from health care and academic organizations, professional associations, the payor and health technology sectors.
- [Primary care panel](#) included experts in chronic non-cancer pain management in the primary care setting, including members of leading health care organizations with ongoing safe prescribing and provider education initiatives.
- [Standards review panel \(SRP\)](#) consisted of representatives from organizations or professional associations who provided a “boots on the ground” point of view and insights into the practical application of the proposed standards.

The prepublication version of the pain assessment and management standards will be available online until the end of 2018. After Jan. 1, 2019, access the standards in the E-dition or standards manual.

### Leadership

**LD.04.03.13: Pain assessment and pain management, including safe opioid prescribing, is identified as an organizational priority by practice leaders.**

### Requirement

LD.04.03.13 EP 1: The practice has a leader or leadership team that is responsible for pain management and opioid prescribing and developing and monitoring performance improvement activities. (See also PI.02.01.01, EP 19)

## Rationale

Gaps in the evidence on optimal pain management, combined with a substantial rise in opioid use and associated harms over the past two decades, are of great concern for health care organizations and the public. Solutions to these safety and quality problems require the coordination of administrative and physician leadership to promote quality initiatives and allocate resources for safe pain management.

## Reference\*

- National Academy of Medicine (NAM). First do no harm: Marshaling clinician leadership to counter the opioid epidemic. Washington, DC: National Academy of Medicine.

## Requirement

LD.04.03.13, EP 3: The practice provides staff and licensed independent practitioners with educational resources to improve pain assessment, pain management, and the safe use of opioid medications based on the identified needs of its patient population.

## Rationale

Provider competence is stressed in clinical practice guidelines, regulations, and by experts in the field. Providers are expected to be knowledgeable about multiple modalities of pain treatment, early identification and prevention of harms of opioid therapy, and management of patients with complex needs. However, providers report insufficient training in pain management and opioid prescribing. The practice can increase provider competence in pain management by providing access to effective education resources.

## Reference\*

- Mostofian F, et al. Changing physician behavior: What works? *The American Journal of Managed Care*. 2015;21(1):75-84.
- Alford DP, et al. SCOPE of pain: An evaluation of an opioid risk evaluation and mitigation strategy continuing education program. *Pain Medicine*. 2016;17(1):52-63.

## Requirement

LD.04.03.13, EP 4: The practice provides information to staff and licensed independent practitioners on available services for consultation and referral of patients with complex pain management needs.

## Rationale

Many patients in hospital and ambulatory settings have complex pain management needs and require multidisciplinary care. Access to specialists through consultation, referral, or use of in-house experts reflects best practice. For example, consultation or referral to a pain specialist is advised when it is necessary to develop a perioperative pain management plan for the patient with opioid tolerance or a history of substance abuse. Cost, lack of transportation, and low availability of services may hinder the use of nonpharmacologic treatments. A barrier also could be the provider's limited knowledge about available networks of nonpharmacologic care providers in the community. Practice leaders can support access by providing information about local resources for patient referral.

## Reference\*

- Chou R, et al. Management of postoperative pain: A clinical practice guideline from the American Pain Society, the American Society of Regional Anesthesia and Pain Medicine, and the American Society of

Anesthesiologists' Committee on Regional Anesthesia, Executive Committee, and Administrative Council. *The Journal of Pain*. 2016;17(2):131-57.

- Vadivelu N, et al. Challenges of pain control and the role of the ambulatory pain specialist in the outpatient surgery setting. *Journal of Pain Research*. 2016;9:425.

## Provision of Care, Treatment, and Services

**PC.01.02.07: The practice assesses and manages the patient's pain and minimizes the risks associated with treatment.**

### Requirement

PC.01.02.07, EP 1: The practice has defined criteria to screen, assess, and reassess pain that are consistent with the patient's age, condition, and ability to understand.

### Rationale

Practices need to develop systems for pain screening and assessment in order to support appropriate individualized pain treatment and perioperative pain management. The practice is responsible for ensuring that appropriate screening and assessment tools are readily available and used appropriately. The tools required to adequately assess pain may differ depending on a patient's age, condition, ability to understand, and whether pain is acute or chronic. For example, chronic pain generally requires more extensive patient assessment, including various domains of physical and functional impairment.

### Reference\*

- Miller RM and Kaiser RS. Psychological characteristics of chronic pain: A review of current evidence and assessment tools to enhance treatment. *Current Pain and Headache Reports*. 2018;22(3):22.
- Reid MC, et al. Management of chronic pain in older adults. *British Medical Journal*. 2015;350(7995):1-0.
- Stanos S, et al. Rethinking chronic pain in a primary care setting. *Postgraduate Medicine*. 2016;128(5):502-15.
- Tobias JD. Acute pain management in infants and children—Part 1: Pain pathways, pain assessment, and outpatient pain management. *Pediatric Annals*. 2014;43(7):e163-8.
- Core Curriculum for Pain Management Nursing, Third Edition. (Eds. M. Czarnecki & H. Turner). Elsevier, 2018.

### Requirement

PC.01.02.07, EP 4: The practice develops a pain treatment plan based on evidence-based practices and the patient's clinical condition, past medical history, and pain management goals.

Note: A pain treatment plan relates to the procedure and treatment provided by the practice.

### Rationale

Differences in the experience of acute or chronic pain may be caused by pain pathophysiology, risk factors, comorbidities, and psychosocial characteristics. These individual variations support an individualized model of pain management, as well as responsible transition of care across care settings to support ongoing care needs. In some instances, an individualized approach to treatment and monitoring is necessary and safe because insufficient clinical guidance exists in areas such as evidence on postoperative opioid tapering protocols.

## Reference\*

- Stanos S, et al. Rethinking chronic pain in a primary care setting. *Postgraduate Medicine*. 2016;128(5):502-15.
- Gordon DB, et al. Research gaps in practice guidelines for acute postoperative pain management in adults: Findings from a review of the evidence for an American Pain Society Clinical Practice Guideline. *The Journal of Pain*. 2016;17(2):158-66.
- Cornelius R, et al. Acute Pain Management in Older Adults. *Journal of Gerontological Nursing*. 2017;43(2):18-27.
- Chou R, et al. Management of postoperative pain: A clinical practice guideline from the American Pain Society, the American Society of Regional Anesthesia and Pain Medicine, and the American Society of Anesthesiologists' Committee on Regional Anesthesia, Executive Committee, and Administrative Council. *The Journal of Pain*. 2016;17(2):131-57.

## Requirement

PC.01.02.07, EP 5: The practice involves patients in the pain management treatment planning process through the following:

- Developing realistic expectations and measurable goals that are understood by the patient for the degree, duration, and reduction of pain
- Discussing the objectives used to evaluate treatment progress (for example, relief of pain and improved physical and psychosocial function)
- Providing education on pain management, treatment options, and safe use of opioid and non-opioid medications when prescribed (See also RI.01.02.01, EPs 2, 4)

## Rationale

Joint Commission's technical advisory panel on pain management emphasized the importance of discussions between the patient and the provider/care team about realistic goals and expectations for the trajectory of pain, especially when the outcomes of pain treatment are uncertain. It is important to identify domains of function or quality of life issues that the patient values, and prioritize improvement in these areas to increase satisfaction with treatment progress.

## Reference\*

- O'Brien EM, et al. Patient-centered perspective on treatment outcomes in chronic pain. *Pain Medicine*. 2010;11(1):6-15.

## Requirement

PC.01.02.07, EP 8: The practice educates the patient and family on discharge plans related to pain management including the following:

- Pain management plan of care
- Side effects of pain management treatment
- If applicable, activities of daily living, including the home environment, that might exacerbate pain or reduce effectiveness of the pain management plan of care, as well as strategies to address these issues
- Safe use, storage, and disposal of opioids when prescribed

## Rationale

Patients perceive good patient-provider communication and education as indicators of high quality care. Discharge education is an opportunity for the provider/care team to engage the patient in a discussion on the pain management plan and opioid safety.

## Reference\*

- Mohammed K, et al. Creating a patient-centered health care delivery system: A systematic review of health care quality from the patient perspective. *American Journal of Medical Quality*. 2016;31(1):12-21.

## Resources

U.S. Food and Drug Administration: [Disposal of unused medicines: What you should know](#)

## Requirement

PC.01.02.07, EP 9: Procedures address the use of intravenous administration of medications and other pain management techniques for patients who experience pain.

**Note: This is a pre-existing requirement (currently EP 5).**

## Performance Improvement

**PI.02.01.01: The practice compiles and analyzes data.**

## Requirement

PI.02.01.01, EP 19: The practice monitors the use of opioids to determine if they are being prescribed safely. (See also LD.04.03.13, EP 1)

## Rationale

Cautious opioid prescribing for chronic and acute pain is promoted in clinical practice guidelines and regulatory policies. Practices can define metrics related to the use of opioids to identify priority areas for practice improvement and resource allocation.

## Reference\*

- Midboe AM, et al. Measurement of adherence to clinical practice guidelines for opioid therapy for chronic pain. *Translational Behavioral Medicine*. 2011;2(1):57-64.
- Parchman ML, et al. Primary care clinic re-design for prescription opioid management. *The Journal of the American Board of Family Medicine*. 2017;30(1):44-51.

\*Not a complete literature review.